Improving Transitions Home for Opioid-Exposed Newborns

Team Hope.

a Vanderbilt initiative to improve the care of pregnant women and infants affected by the opioid crisis, sought to improve transitions home from the hospital after birth.

Their discharge checklist:

Follow-up appointment with primary care physician scheduled prior to discharge

Referral to home visiting services

Referral to Early Intervention Services

If diagnosed with neonatal abstinence syndrome, referral to Pediatric Development Clinic

If exposed to hepatitis C virus, referral for follow-up and testing

Opioid-exposed newborns, especially those diagnosed with neonatal abstinence syndrome, are at risk of poor outcomes after discharge from the hospital. including:



Hospital Readmission



Emergency Department Utilization



Adverse Developmental **Outcomes**

Newborns connected to discharge services:

before checklis

after checklis

Implementation of a checklist improves the connection to important follow-up services for moms and babies.

Crook, TW, Munn, EK, Scott, TA, Morad, A, Wyatt, J, Johnson, DP, White, M, Patrick, SW. Improving the Discharge Process for Opioid Exposed Neonates. Hospital Pediatrics, 2019, DOI: 10.1542/hpeds.2019-0088 Supported by The Memorial Foundation and the Boedecker Foundation.

www.childpolicy.org • @VUMCchildpolicy • For more info, contact teamhope@vumc.org

